

EFFECT OF NPK, MICRONUTRIENTS AND N-PLACEMENT ON THE GROWTH AND YIELD OF SUNFLOWER

MUZZAMMIL HUSSAIN SIDDIQUI*, FATEH CHAND OAD*,
M. KALEEM ABBASI** and ALLAH WADHAYO GANDAH*^I

* Sindh Agriculture University, Tandojam, Pakistan

** Faculty of Agriculture, Rawalakot, AJK, Pakistan

ABSTRACT

A series of experiments was carried out at Students Farm, Sindh Agriculture University, Tandojam to evaluate the effect of NPK, micronutrients and N-placement on the growth and yield of sunflower during 2005-2006. The sunflower cv. HO-1 was studied under 7 different NPK levels and placement methods (0-0-0 NPK kg ha⁻¹, 60-30-30 NPK kg ha⁻¹ (N as broadcast), 60-30-30 NPK kg ha⁻¹ (N as fertigation), 90-45-45 NPK kg ha⁻¹ (N as broadcast), 90-45-45 NPK kg ha⁻¹ (N as fertigation), 120-60-60 NPK kg ha⁻¹ (N as broadcast), 120-60-60 NPK kg ha⁻¹ (N as fertigation) and 7 varying zinc and boron doses (0-0, 10.0-1.5, 10.0-2.0, 15.0-1.5, 15.0-2.0, 20.0-1.5 and 20.0-2.0 Zn-B kg ha⁻¹ respectively). Both macronutrient and micronutrient combinations showed significant differences and enhanced plant parameters. The values of all crop parameters increased as the dose of fertilizers increased. The tallest plant height, maximum stem girth, largest head diameter, most seeds head⁻¹, heaviest seed weight head⁻¹, largest seed index and best seed yield (1738.0 kg ha⁻¹) were recorded under application of 90-45-45 NPK with 15-1.5 Zn-B kg ha⁻¹ (N applied as fertigation). Beyond these level of macro- and micronutrients responses were not significant. The traditional fertility application, broadcast N at 120-60-60 NPK with 15-1.5 Zn-B kg ha⁻¹ was no more effective in achieving higher values of crop parameters than comparable N-fertigated plots. The reason for increased values in N-fertigated plots may be that these plots received equal amounts of N in whole the plot during irrigation as compared to broadcast method. This study proved the incorporation of 90-45-45 kg ha⁻¹ (N as fertigation) with 15-1.5 Zn-B kg ha⁻¹ to be the optimum fertilizer dose for sunflower growth and yield. This could save growers on costs for NPK.

Keywords: Sunflower, Nitrogen, Phosphorus, Potassium, Micronutrients, Fertigation, Placement, Growth, Yield

Citation: Siddiqui, M.H., F.C. Oad, M.K. Abbasi and A.W. Gandahi. 2009. Effect of NPK, micronutrients and N-placement on the growth and yield of sunflower. Sarhad J. Agric. 25(1): 45-52.

INTRODUCTION

Sunflower is the fourth oil grain crop grown worldwide by area (Fagundes *et al.*, 2007). Fertilizer management is an important part of crop production. This is due in part to increasing costs for fertilizer materials, particularly nitrogen (N) fertilizer. Producers need to know how different production inputs affect crop yields so that they can choose technically efficient input combinations. They also need to know how input usage affects the profitability of their operations with varying input and crop prices. Fertilizer is an input that can easily be reduced, thus lowering input costs. However, fertilizer is a crop input that has a high return on investment if applied at appropriate level. Prediction of optimum fertilizer rates is important because of increasing economic and environmental concerns. The economic optimum N rate was approximately 110 kg ha⁻¹ for sunflower grown under irrigated conditions of Anatolia, Turkey (Peker and Ozer, 2005).

Numerous studies have shown that NO₃-N leaching is a common and sometimes serious problem when inorganic N and organic wastes are used (Roth and Fox, 1990; Weil *et al.* 1990; Schepers *et al.*, 1991; Spalding and Exner, 1993; Bacon, 1995; Moreno *et al.*, 1996). A survey of groundwater quality in China indicates that the nitrate level of groundwater is higher in intensively cropped agricultural regions where the primary crops are vegetables and where high rates of water and nitrogen are applied, than in regions where grains are the primary crops (Ju and Zhang, 2003). Leaching of NO₃-N is economically and environmentally undesirable (Ryden *et al.*, 1984; Katyal *et al.*, 1985; Poss and Saragoni, 1992; Jones and Schwab, 1993; Theocharopoulos *et al.*, 1993). The US Environmental Protection Agency (EPA, 1990) has set 10 mg L⁻¹ N as the NO₃-N drinking water standard as a safeguard against methemoglobinemia from developing in infants (Fletcher, 1991). Additionally, groundwater with high nitrate levels that discharge into sensitive surface waters can contribute to the long-term eutrophication of these

water bodies. Combined application of irrigation and nitrogen through fertigation is now becoming a common practice in modern agriculture because of its advantages over broadcast N application. Some of these advantages [(Gascho *et al.* 1984; Threadgill, 1985; Criado, 1996; Hagin and Lowengart, 1996)] are more timely nitrogen application, more uniform nitrogen application, reduced environmental contamination, faster movement of applied N into the rooting zone and lessened soil compaction and crop damage related to mechanical side dressing. Concerns about best management practice are increasing because mismanagement of nitrogenous fertilizers can lead to contamination of ground and surface water sources as well as soil degradation. In this regard fertigation had a consistent effect on total nitrogen use efficiency (Hou *et al.*, 2007) and a more uniform initial NO₃-N distribution produced a more uniform NO₃-N distribution through fertigation (Jiusheng *et al.*, 2005). Similar observations of increased uptake as a result of fertigation were reported earlier by Vasane *et al.* (1996). Alva and Mozzafari (1995) also reported that fertigation treatments maintained high concentration of NO₃-N at shallower depth than did broadcast treatments. Further, it was also reported that reduced loss of nutrients could be through fertigation compared to soil application of fertilizer (Hebbar *et al.*, 2004).

There are positive effects of micronutrient application on the growth of sunflower, in terms of plant height, number of leaves and dry matter production per plant. These effects can be contributed to enhanced metabolic function of micronutrients in the plant (Amberger, 1980). Under Egyptian conditions, positive effects of micronutrients on crops were reported in several studies by El-Fouly *et al.* (1990) and Abdalla and Mobarak (1992) found that increases in plant growth in terms of plant height, dry weight and uptake due to continuous addition of Fe, Mn and Zn through irrigation water. Sunflower is also one of the most sensitive crops to low boron supply and deficiency. Among the micronutrients, zinc deficiency is most widespread on a wide range of soils (Graham *et al.*, 1992; Cakmak *et al.*, 1996; Grewal *et al.*, 1997). Incorporation of Zn with other fertilizers into the normal cultivation zone of the soil (top 10 cm) is a common practice to overcome the Zn deficiency problem. Because Zn is readily fixed in soil, little of this Zn moves down the profile below the cultivation zone (Brennan and McGrath, 1988). The role of Zn in subsoil nutrition is of particular interest because of its importance in maintaining membrane integrity of root cells (Cakmak and Marschner, 1988; Welch, 1995) and the possible role of zinc in reducing the toxic effects of excessive B (Graham *et al.*, 1987; Singh *et al.*, 1990); excessive B in subsoil is a problem. Sunflower roots are also sensitive to B deficiency as they stop their growth in <6 h after the removal of B from the growth medium (Dugger, 1983). The recent observations demonstrate that the concentration of B in soils and plants not only varies with soil type, plant species and environmental conditions, but also its excess or deficiency may affect the plant growth and production, because there is small concentration range between deficiency and toxicity in soil-plant system (Tariq and Mott, 2007). Thus, application of adequate micronutrient fertility increased crop yields, improved the nutrient element concentrations in plant tissue, and enhanced soil macro and micronutrient status (Adediran *et al.*, 2004) and can give 67% more yield over control (Taiwo *et al.* 2001). Looking at the above considerations, the study was conducted to explore appropriate rates and placement methods of macro- and micronutrient for sunflower production.

MATERIALS AND METHODS

This field research was conducted at Sindh Agriculture University, Tandojam, Pakistan, during spring and autumn seasons in both 2005 and 2006. The experiment was set in a randomized complete block design (factorial) having three replications. The soil was clay loam in texture, non saline (EC 0.96 dS m⁻¹), alkaline in reaction (pH 8.4), calcareous (CaCO₃ 9.8%), low in organic matter (0.57%), available phosphorus (3.06 mg kg⁻¹) and total nitrogen content was 0.05%.

Treatments

Factor A

The NPK treatments studied were:

(0-0-0 NPK kg ha⁻¹, 60-30-30 NPK kg ha⁻¹ (N as broadcast), 60-30-30 NPK kg ha⁻¹ (N as fertigation), 90-45-45 NPK kg ha⁻¹ (N as broadcast), 90-45-45 NPK kg ha⁻¹ (N as fertigation), 120-60-60 NPK kg ha⁻¹ (N as broadcast) and 120-60-60 NPK kg ha⁻¹ (N as fertigation).

Factor B

The micronutrient treatments studied were:

Different zinc and boron doses (0-0, 10.0-1.5, 10.0-2.0, 15.0-1.5, 15.0-2.0, 20.0-1.5 and 20.0-2.0 Zn-B kg ha⁻¹, respectively).

The sunflower variety HO-1 was sown in rows 75 cm apart. Plants were thinned to a spacing of 25 cm w/c rows. The crop was cleared from weeds by one hoeing with Spade before the 1st irrigation. All other cultural practices were adopted uniformly in all the plots to maintain the experimental area.

Fertilizer Placement Methodology

Broadcast

This is practice generally used by farmers. In this method, N fertilizer was incorporated as urea on the surface of soil under three split applications i.e., at the same time as the first, second and third irrigations. However, all the P from triple super phosphate (TSP) and K from muriate of potash (MOP) were applied as indicated above for all treatments during seed bed preparation.

Fertigation

In this method urea was applied in three splits during first, second and third irrigations as fertigation (application of fertilizer with irrigation water). The appropriate urea rate was mixed with water in a tank. The tank was then placed on the head of channel which released droplets in the running water pertaining to the appropriate plots.

Statistical Analysis

Data recorded individually for each parameter were subjected to the ANOVA technique by using MSTATC computer software and the significant means were separated by using Fisher's Protected LSD test (Steel and Torrie, 1980).

RESULTS AND DISCUSSION

The experimental results revealed that all macro- and micronutrient treatments significantly enhanced plant height, stem girth, head diameter, seeds head⁻¹, seed weight head⁻¹, seed index, and seed yield of sunflower crop. Further, as NPK doses increased from low (60-30-30 NPK kg ha⁻¹) to high (120-60-60 NPK kg ha⁻¹), significant differences in growth parameters were noted. This was also true for micronutrients (Zn and B), where values of all crop parameters increased as the dose of micronutrients increased. Taller plants (202.20 cm), greater stem girth (12.26 cm), larger head diameter (23.04 cm), more seeds head⁻¹ (696.46), heavier seed weight head⁻¹ (49.06 g), greater seed index (71.28 g), satisfactory seed yield (2743.07 kg ha⁻¹) and higher dry matter (11666.70 kg ha⁻¹) were recorded under application of 90-45-45 NPK kg ha⁻¹ with 15 kg ha⁻¹ Zn and 1.5 kg ha⁻¹ B (N applied as fertigation). Statistically, beyond this level, non-significant increases in plant parameters studied were exhibited (Table I-VII). The traditional broadcast N application at higher doses (120-60-60 NPK with 15 kg ha⁻¹ Zn and 1.5 kg ha⁻¹ B) did not result in recording higher values of crop parameters as compared to N-fertigated plots. The seed yield of sunflower showed increasing response to traditional application of higher fertilizer rates up to 120-60-60 NPK kg ha⁻¹ but the values were low as compared to 90-45-45 NPK kg ha⁻¹ where N applied as fertigation. This may be the reason that N-fertigated plots received equal amounts of N in whole the plot where N might have distributed uniformly in the field as compared to broadcast method. Recently, the farmers are using 120-60-60 NPK kg ha⁻¹ as broadcast, but with N-fertigation, the reduced application of 90-45-45 kg ha⁻¹ was found to result in equivalent yield with lower fertilizer costs. The present results deviate from the previous study conducted by Ali *et al.* (2004) which showed broadcast application of 150 kg N ha⁻¹ increased plant height, head diameter, number of filled seeds per head, 1000-seed weight, biological and seed yields and seed oil content. Malik *et al.* (2004) also observed that increasing levels of N, P and K increased yield and yield components of sunflower (plant height, head diameter, number of achenes head⁻¹, 1000-achene weight) but decreased the seed oil content. Their seed yield (1231.47 kg ha⁻¹) was recorded with broadcast application of 130-90-90 NPK kg ha⁻¹. Nawaz *et al.* (2003) found the optimum fertilizer requirement as broadcast of 120-90-60 NPK kg ha⁻¹ for sunflower crop. The higher nitrogen broadcast dose of 160 kg N ha⁻¹ was exhibited by Ali *et al.* (2000) for maximum seed yield per head by sunflower plants. Other researchers from Pakistan also suggested higher fertilizer doses as broadcast for sunflower crop. Farid *et al.* (2000) recommended 180-120 kg NP ha⁻¹; Hussain *et al.* (1998) observed 125 kg N ha⁻¹; Bakht *et al.* (1989) suggested 120 kg N ha⁻¹. Further, Ali *et al.* (1997) also stated that plant height, stover yield and number of seeds per floral disc increased with N application, with 100 kg N ha⁻¹ the optimum rate. Their yield and yield components were not significantly different between P application methods (50 kg P₂O₅ ha⁻¹ applied by broadcasting or fertigation).

Hussain *et al.* (1998) reported that yield increased with up to 125 kg N ha⁻¹. Meo *et al.* (1999) found that dry weights of leaves, stems, roots and seeds decreased with decreasing N rate and increasing water stress. Farid *et al.* (2000) also recommended higher doses for sunflower production and reported that application of 180 kg N and 120 kg P₂O₅ ha⁻¹ gave the highest yield of 1249.2 kg ha⁻¹ of sunflower seed while control plots only yielded 384.3 kg ha⁻¹.

In this study, among the N-placement methods, the N-fertigation was found efficient for growth and yield of sunflower. Cadahia (1993) reported that the advantages of fertigation showed a slow-release fertilizer and lower loss of N and therefore a lower degree of ground water contamination which not only increased the N uptake by the plant as well as the leaf and root weight, but also produced higher yields. This technique can reduce fertilizer application costs by eliminating an operation and improving nutrient efficiency. Also, it could conceivably reduce leaching or denitrification (gaseous) losses of nitrogen and lower the luxury uptake of nutrients by plants. Fertigation enables users to put the fertilizers in the plant root zone or on the canopy at the desired frequency, amount and concentration at appropriate time (Kumar *et al.*, 2000).

In this study, the incorporation of Zn and B as micronutrients also enhanced growth and seed yield of sunflower and 15 kg ha⁻¹ Zn and 1.5 kg ha⁻¹ B was recorded as optimum dose with N-fertigated plots (Table I-VII). The response of Zn and B in sunflower might be due to the reason that the site was deficient in both these micronutrients, and previous reports also supports that soils of Asia have shortfall in the availability of micronutrients (Asia Feb., 1998). Gitte *et al.* (2005) was in the view that combined application of Zn (5.25 kg ha⁻¹) and B (6.3 kg ha⁻¹) was more effective than Zn (5.25 kg ha⁻¹) alone or B (2.1 kg ha⁻¹) alone. The combined application of Zn and B exhibited increases over the control in straw yield, harvest index, translocation efficiency, 100-seed weight and seed yield. Meanwhile, Marchetti *et al.* (2001) found borax at 1.0 mg/dm³ to exhibit the highest yield. Castro *et al.* (2002) suggested that boron could also be used with herbicides and application of B with herbicides is a good strategy for avoiding B deficiency while controlling weeds in sunflower. Likewise, Vyakaranahal *et al.* (2001) reported that B at 0.1% spray at ray floret stage increased the processed seed yield by 49.0 and 43.1% in kharif and summer sunflowers, respectively. The foliar sprays also increased the capitulum diameter, number of filled seeds per capitulum, seed set percentage, seed yield per plant, 100 seed weight, volume weight of seed, oil content in seed, germination percentage, seedling dry weight and seedling vigor index. Reddy and Shaik (2000) conducted experiments on various fertilizers and reported that 100% recommended fertilizer dose +500 kg gypsum ha⁻¹ + 0.2% borax gave the highest net returns and cost:benefit ratio as a result of increased yield due to boron application. Further, Renukadevi *et al.* (2003) reported that boron nutrition increased the head diameter, number of filled seeds and 100-seed weight (5.8 g).

Table I. Plant height (cm) of sunflower as affected by NPK, Zn, B and placement methods

Zinc + Boron (kg ha ⁻¹)	NPK (kg ha ⁻¹)	60-30-30 broadcast	90-45-45 broadcast	120-60-60 broadcast	60-30-30 N-fertigation	90-45-45 N-fertigation	120-60-60 N-fertigation
0-0	69.79 m	89.73 l	114.65 ij	140.59 f	111.83 j	142.57 ef	145.21 def
10-1.5	70.76 m	100.63 k	127.61 gh	155.55 cd	122.62 hi	158.53 c	161.16 c
10-2.0	72.90 m	110.67 j	139.58 f	185.45 b	137.25 fg	181.43 b	189.06 b
15-1.5	74.82 m	124.63 h	152.05 cde	206.73 a	150.88 cde	207.20 a	209.48 a
15-2.0	74.14 m	125.12 h	152.53 cde	207.39 a	151.39 cde	207.45 a	209.81 a
20-1.5	74.54 m	126.61 h	154.53 cd	208.30 a	153.54 cd	207.70 a	210.07 a
20-2.0	75.60 m	128.11 gh	155.53 cd	209.71 a	155.40 cd	208.29 a	210.57 a

SE= 2.485, LSD(5%) = 6.931, LSD(1%)= 9.143

Table II. Stem girth (cm) of sunflower as affected by NPK, Zn, B and placement methods

Zinc + Boron (kg ha ⁻¹)	NPK (kg ha ⁻¹)							
	0-0-0	60-30-30 broadcast	90-45-45 broadcast	120-60-60 broadcast	60-30-30 N-fertigation	90-45-45 N-fertigation	N- fertigation	120-60-60 N- fertigation
0-0	5.08 m	7.17 l	8.75 ghij	9.60 defg	8.51 hijk	9.37 defgh		9.85 def
10-1.5	5.17 m	7.63 kl	9.03 fgghi	10.34 d	9.06 fgghi	10.21 de		10.36 d
10-2.0	5.23 m	7.92 jkl	9.29 efgh	11.40 bc	9.07 fgghi	11.29 c		11.44 bc
15-1.5	5.35 m	8.11 ijk	10.06 de	12.30 ab	9.63 defg	12.26 ab		12.410 a
15-2.0	5.39 m	8.17 ijk	10.07 de	12.42 a	9.74 def	12.32 ab		12.537 a
20-1.5	5.39 m	8.18 ijk	10.24 de	12.52 a	9.84 def	12.42 a		12.588 a
20-2.0	5.39 m	8.19 ijk	10.08 de	12.63 a	9.94 def	12.52 a		12.708 a

SE = 0.2309, LSD(5%)= 0.6441, LSD(1%) = 0.8496

Table III. Head diameter (cm) of sunflower as affected by NPK, Zn, B and placement methods

Zinc + Boron (kg ha ⁻¹)	NPK (kg ha ⁻¹)							
	0-0-0	60-30-30 broadcast	90-45-45 broadcast	120-60-60 broadcast	60-30-30 N-fertigation	90-45-45 N-fertigation	N- fertigation	120-60-60 N- fertigation
0-0	7.00 l	8.78 l	12.49 k	14.99 ghi	11.43 k	15.34 fgh		15.44 fgh
10-1.5	7.57 l	11.17 k	15.69 fgh	19.91 bc	13.10 ijk	19.11 bcd		19.29 bc
10-2.0	7.46 l	11.59 k	17.15 def	20.61 b	14.66 hij	20.82 b		20.90 b
15-1.5	7.56 l	12.94 jk	18.38 cde	22.72 a	16.73 efg	23.04 a		23.15 a
15-2.0	7.57 l	12.96 jk	18.40 cde	23.26 a	16.76 efg	23.10 a		23.22 a
20-1.5	7.43 l	12.98 jk	18.42 cde	23.31 a	16.78 efg	23.14 a		23.27 a
20-2.0	7.94 l	13.00 jk	18.43 cde	23.37 a	16.80 efg	23.19 a		23.31 a

SE= 0.4948, LSD(5%)= 1.380, LSD(1%)= 1.820

Table IV. Seeds head¹ of sunflower as affected by NPK, Zn, B and placement methods

Zinc + Boron (kg ha ⁻¹)	NPK (kg ha ⁻¹)							
	0-0-0	60-30-30 broadcast	90-45-45 broadcast	120-60-60 broadcast	60-30-30 N-fertigation	90-45-45 N-fertigation	N- fertigation	120-60-60 N- fertigation
0-0	378.71 k	543.81 j	553.77 ij	591.22 efg	554.56 ij	590.94 efg		608.19 bcdef
10-1.5	389.39 k	554.22 ij	576.71 ghi	599.27 defg	566.43 hij	598.51 defg		599.78 defg
10-2.0	388.52 k	562.40 ij	590.37 fg	613.20 bcdef	589.36 fgh	602.80 cdef		614.05 bcdef
15-1.5	395.71 k	617.21 bcd	630.18 b	696.57 a	629.73 b	696.46 a		696.74 a
15-2.0	397.99 k	616.32 bcde	629.05 b	695.36 a	628.17 bc	695.24 a		696.39 a
20-1.5	400.10 k	616.44 bcde	629.18 b	695.47 a	628.30 bc	695.67 a		696.56 a
20-2.0	401.18 k	616.63 bcde	629.27 b	695.58 a	628.38 bc	695.43 a		715.57 a

SE=5.985, LSD(5%)= 16.69, LSD(1%)= 22.02

Table V. Seeds weight head¹ (g) of sunflower as affected by NPK, Zn, B and placement methods

Zinc + Boron (kg ha ⁻¹)	NPK (kg ha ⁻¹)							
	0-0-0	60-30-30 broadcast	90-45-45 broadcast	120-60-60 broadcast	60-30-30 N-fertigation	90-45-45 N-fertigation	N- fertigation	120-60-60 N- fertigation
0-0	14.71 i	27.92 h	31.50 g	35.83 ef	31.52 g	35.75 ef		36.90 cde
10-1.5	15.14 i	32.52 g	34.06 fg	37.94 bcde	33.99 fg	37.88 bcde		38.03 bcde
10-2.0	15.11 i	34.05 fg	36.24 def	39.79 bc	35.58 ef	39.04 bcd		39.89 b
15-1.5	15.43 i	39.00 bcd	40.54 b	49.62 a	40.55 b	49.06 a		49.64 a
15-2.0	15.53 i	38.96 bcd	40.58 b	49.60 a	40.47 b	49.56 a		49.73 a
20-1.5	15.63 i	38.97 bcd	40.60 b	49.59 a	40.48 b	49.60 a		49.75 a
20-2.0	15.67 i	38.98 bcd	40.60 b	49.59 a	40.48 b	49.59 a		51.30 a

SE = 0.685, LSD(5%)= 1.913, LSD(1%)= 2.424

Table VI. Seed Index (g) of sunflower as affected by NPK, Zn, B and placement methods (Mean of two years)

Zinc + Boron (kg ha ⁻¹)	NPK (kg ha ⁻¹)		90-45-45 broadcast	120-60-60 broadcast	60-30-30 N-fertigation	90-45-45 N-fertigation	120-60-60 N-fertigation
	0-0-0	60-30-30 broadcast					
0-0	38.84 g	55.76 f	56.78 ef	60.62 cd	56.86 ef	60.59 cd	60.65 cd
10-1.5	38.88 g	58.69 de	59.00 de	63.31 bc	60.01 d	63.30 bc	63.41 bc
10-2.0	38.90 g	59.66 de	60.53 cd	64.88 b	60.43 cd	64.76 b	64.96 b
15-1.5	39.03 g	63.20 bc	64.48 b	71.29 a	64.40 b	71.28 a	71.34 a
15-2.0	39.04 g	63.19 bc	64.50 b	71.30 a	64.41 b	71.29 a	71.41 a
20-1.5	39.08 g	63.21 bc	64.51 b	71.31 a	64.42 b	71.30 a	71.43 a
20-2.0	39.08 g	63.23 bc	64.52 b	71.32 a	64.43 b	71.31 a	71.67 a

SE = 0.7404, LSD(5%)= 2.065, LSD(1%)= 2.724

Table VII. Seed yield (kg ha⁻¹) of sunflower as affected by NPK, Zn, B and placement methods

Zinc + Boron (kg ha ⁻¹)	NPK (kg ha ⁻¹)		90-45-45 broadcast	120-60-60 broadcast	60-30-30 N-fertigation	90-45-45 N-fertigation	120-60-60 N-fertigation
	0-0-0	60-30-30 broadcast					
0-0	815.70 i	1548.56 h	1746.85 g	1987.31 cdef	1914.62 efg	1985.39 cdef	1996.94 cdef
10-1.5	839.90 i	1803.71 g	1888.77 fg	2086.09 h	1885.17 fg	2101.37 bcd	2114.70 bcd
10-2.0	838.04 i	1888.66 fg	1981.98 def	2262.13 b	2001.01 cdef	2164.74 bc	2267.50 b
15-1.5	855.79 i	2163.0bcd	2247.45 b	2746.26 a	2248.81 b	2743.07 a	2752.77 a
15-2.0	861.61 i	2159.9bcd	2250.72 b	2750.36 a	2247.03 b	2749.45 a	2757.65 a
20-1.5	866.90 i	2161.1bcd	2251.67 b	2752.86 a	2248.78 b	2751.97 a	2759.09 a
20-2.0	869.06 i	2162.03bcd	2251.34 b	2755.66 a	2249.09 b	2754.60 a	2789.34 a

SE= 42.10, LSD(5%)= 117.40, LSD(1%)= 154.90

CONCLUSION

The application of NPK macronutrient and Zn and B micronutrient fertility significantly enhanced yield and yield contributing parameters of sunflower. The values of all crop parameters increased as the dose of fertilizers increased and maximum response was exhibited under 90-45-45 NPK with 15 kg ha⁻¹ Zn and 1.5 kg ha⁻¹ B (with N applied as fertigation). Beyond this fertility level, non-significant responses were noted. Hence, it may be concluded that application of this treatment seems to be most beneficial for sunflower production.

REFERENCES

- Abdalla, F. E. and Z. M. Mobarak. 1992. Shoot intake of nutrients from different micronutrients fertilizer formulations in faba bean. *African J. Agric. Sci.* 19:147-160.
- Adediran, J. A., L. B. Taiwo, M. O. Akande, O. J. Idowu, R. A. Sobulo, and J. A. Adediran. 2004. Application of organic and inorganic fertilizer for sustainable maize and cowpea yields in Nigeria. *J. Plant. Nut.* 27(7):1163-1181.
- Ali, C.K., M.A. Nadeem and M.A. Javid. 1997. Effect of nitrogen levels and P application strategy on sunflower. 1997. *Fertilizer News.* 42(3):43-47.
- Ali, H., H.A. Randhawa and M. Yousaf. 2004. Quantitative and qualitative traits of sunflower (*Helianthus annuus* L.) as influenced by planting dates and nitrogen application. *Int. J. Agric. and Biol.* 6(2):410-412.
- Ali, M., S.K. Khalil and K. Nawab. 2000. Response of sunflower hybrids to various levels of nitrogen and phosphorus. *Sarhad J. Agric.* 16(5):477-483.
- Alva, A. K. and M. Mozzafari. 1995. Nitrate leaching in a deep sandy soil as influenced by dry broadcast or fertigation of nitrogen for citrus production. In: *Int. Symp. on Fertigation Technology, Israel.* 26-31 March 1995, pp. 67-77.
- Amberger, A. 1980. Foliar application of micronutrients, uptake and incorporation into metabolism. 2nd Workshop 'Micronutrients and Plant Nutrition', Mariut, Egypt, 1979, NRC, Cairo, Egypt. pp. 47-60.
- Asia Feb. 1998. Countering the yield ceiling: A key ingredient in balanced nutrition, *Asia.* 21:36-38.
- Bacon, P. E. 1995. Nitrogen fertilization in the environment, 1st Ed. Marcel Dekker, New York. 608 p.
- Bakht, J., S.K. Khalil, Z. Shah, Ehsanullah and A. Qayyum. 1989. Yield and yield components of maize and sunflower sown alone and in different combination under various levels of nitrogen. *Sarhad J. Agric.* 5(2):101-106.
- Brennan, R. F. and J.F. McGrath. 1988. The vertical movement of zinc on sandy soils in southern Western Australia. *Aust. J. Agric. Res.* 26:211-216.
- Cadahia, C. 1993. Pre-plant slow-release fertilization of strawberry plants before fertigation. *Fert. Res.* 34(3):191-195.
- Cakmak, I., and H. Marschner. 1988. Increase in membrane permeability and exudation in roots of zinc deficient plants. *J. Plant Physiol.* 132:356-361.

- Cakmak, I., A. Yilmaz, M. Kalayci, H. Ekiz, B. Torun, B. Erenoglu and H. J. Braun. 1996. Zinc deficiency as a critical problem in wheat production in central Anatolia. *Plant Soil*. 180:165-172.
- Castro, C., A. M. Brighenti and A. Oliveira Jr. 2002. Tank-mix of boron and herbicides on conventional sunflower sowing. *Planta Daninha*. 20(1): 83-91.
- Criado, S. R. 1996. Considerations on main factors which take part in nitrate contamination of groundwater in Spain with relationship to other EU countries. *Fert. Res.* 43:203-207.
- Dugger W. M. 1983. Boron in plant metabolism. *Encyclopedia of plant physiology, inorganic plant nutrition*, 15B. New York. Springer Verlag. 626-650.
- El-Fouly, M. M., A. F. A.Fawzi, Z. M. Mobarak, E. A. Aly and F. E. Abdalla. 1990. Micronutrient foliar intake by different crop plants, as affected by accompanying urea. *Plant Nutrition Physiology and Applications*. Kluwer Acad. Pub., London. pp. 267-273.
- Environmental Protection Agency. 1990. National groundwater quality survey. Office of Drinking Water, Washington, DC. 211p.
- Fagundes, J., D. Santiago, G. Mello, A. M. deBellé, and N.A. Streck, N. A. 2007. Growth, development and delay of leaf senescence in pot-grown sunflower (*Helianthus annuus* L.): sources and rates of nitrogen. *Ciência Rural*. 37(4):987-993.
- Farid, S. S., Z. Hussain, S. Farida and S. Khan. 2000. Performance of sunflower intercropped in ratoon sugarcane under NP fertilizers. *Sarhad J. Agric.* 16(4):431-434.
- Fletcher, D. A. 1991. A national perspective. *Managing Nitrogen for Groundwater Quality and Farm Profitability*. ASA, SSSA, Madison, WI. pp. 9-17.
- Gascho, G. J., J. E. Hook and G. A. Mitchell. 1984. Sprinkler-applied and side-dressed nitrogen for irrigated corn grown on sand. *Agron. J.* 76:77-81.
- Gitte, A. N., S. R. Patil and M. A. Tike. 2005. Influence of zinc and boron on biochemical and yield characteristics of sunflower. *Indian J. Plant Physio.* 10 (4): 400-403.
- Graham, R. D, R. M. Welch, D. L. Grunes, E. E. Cary and W. A. Norvell. 1987. Effect of zinc deficiency on the accumulation of boron and other mineral nutrients in barley. *Soil Sci. Soc. Amer. J.* 51:652-657.
- Graham, R. D., J. S. Ascher and S. C. Hynes. 1992. Selecting zinc-efficient genotypes for soils of low zinc status. *Plant Soil*. 146:241-250.
- Grewal, H. S., Z. Lu and R. D. Graham. 1997. Influence of subsoil zinc on dry matter production, seed yield and distribution of zinc in oilseed rape genotypes differing in zinc efficiency. *Plant Soil*. 192:181-189.
- Hagin, J., and A. Lowengart. 1996. Fertigation for minimizing environmental pollution by fertilizers. *Fert. Res.* 43:5-7.
- Hebbbar, S. S., B. K. Ramachandrappa, H. V. Nanjappa, and M. Prabhakar. 2004. Studies on NPK drip fertigation in field grown tomato (*Lycopersicon esculentum* Mill.). *European J. Agron.* 21(1):117:127.
- Hou, Z., Li. Pinfang, Li. Baoguo, Gong. Jiang, Wang, and Yanna. 2007. Effects of fertigation scheme on N uptake and N use efficiency in cotton. *Plant Soil*. 290(1-2):115-126.
- Hussain, G., Amanullah, G. Hassan, and A. Rashid. 1998. Response of sunflower cultivars to different nitrogen levels under D.I. Khan conditions. *Sarhad J. Agric.* 14(5):411-415.
- Jiusheng, Li, Bei Li, and M. Rao. 2005. Spatial and temporal distributions of nitrogen and crop yield as affected by non-uniformity of sprinkler fertigation. *Agric. Water Manag.* 76(3):160-180.
- Jones, R. D. and A. P. Schwab. 1993. Nitrate leaching and nitrate occurrence in a fine-textured soil. *Soil Sci.* 155:272-282.
- Ju, X. and F. Zhang. 2003. Nitrate accumulation and its implications to environment in north China. *Ecol. Envir.* 12(1):24-28.
- Katyal, J. C., B. Singh, P. L. G. Vlek and E. T. Crasweel. 1985. Fate and efficiency of nitrogen fertilizers applied to wetland rice II. Punjab, India. *Fertil. Res.* 6:279-290.
- Kumar, S., R. Asrey and R.S. Yojana. 2000. Fertigation: Need of modern agriculture. *Fert. Res.* 44:31,32,37.
- Malik, M.A., M.F. Saleem, M. Sana, and A. Rehman. 2004. Suitable level of N, P and K for harvesting the maximum economic returns of sunflower (*Helianthus annuus* L.). *Int. J. Agric. Biol.* 6(2):240-242.
- Marchetti, M. E., W. R. Motomya, A. C. Fabrício and J. O. Novelino. 2001. Sunflower, *Helianthus annuus*, response to sources and levels of boron. *Acta Scientiarum.* 23 (5): 1107-1110.
- Meo, A.A., F. Baig, Z. Khan, and W. Naseem. 1999. Effect of urea and sporadic drought on dry matter and yield of sunflower (*Helianthus annuus* L.). *Sarhad J. Agric.* 15(5):443-446.
- Moreno, F., J. A. Cayuela, J. E. Fernandez, E. Fernandez-Boy, J. M. Murillo and F. Cabrera. 1996. Water balance and nitrate leaching in an irrigated maize crop in SW Spain. *Agric. Water Manage.* 32:71-83.
- Nawaz, N., G. Sarwar, M. Yousaf, T. Naseeb, A. Ahmed and M.J. Shah. 2003. Yield and yield components of sunflower as affected by various NPK levels. *Asian J. Plant Sci.* 2(7):561-562.
- Peker, K. and H. Ozer. 2005. Determination of economically optimal rate of nitrogen fertilizer for irrigated sunflower in Eastern Anatolia, Turkey. *Bulgarian J. Agric. Sci.* 11(3):323-330.
- Poss, R. and H. Saragoni. 1992. Leaching of nitrate, calcium and magnesium under maize cultivation on an oxisol in Togo. *Fert. Res.* 33:123-133.
- Reddy, M.P. and M. Shaik. 2000. Influence of nitrogen and phosphatic fertilizer on growth, yield components and yield of sunflower. *Crop Res.* 20(2):293-296.

- Renukadevi, A., P. Savithri and K. Andi. 2003. Sources, levels and methods of micronutrients application on the dry matter production, yield attributes and yield of sunflower *Helianthus annuus* crop. *Crop Res.* 25(3):436-440.
- Roth, L.V. and R.H. Fox. 1990. Soil nitrate accumulations following nitrogen-fertilized corn in Pennsylvania. *J. Envir. quality.* 19:243-248.
- Ryden, J. C., P.R. Ball and E.A. Garwood. 1984. Nitrate leaching from grassland. *Nature.* 311:50-53.
- Schepers, J.S., M. G. Moravek, E. E. Alberts, and K. D. Frank. 1991. Maize production impacts on groundwater quality. *J. Envir. Quality.* 20:12-16.
- Singh, J.P., D.J. Dahiya and R P. Narwal. 1990. Boron uptake and toxicity in wheat in relation to zinc supply. *Fert. Res.* 24:105-110.
- Spalding, R.F. and M.E. Exner. 1993. Occurrence of nitrate in groundwater, A Review. *J. Envir. Quality.* 22:392-402.
- Steel, R.G.D. and J.H. Torrie. 1980. Principles and procedures of statistics. McGraw. Hill Book Co. Inc. NewYork. 481p.
- Taiwo, L.B., J.A. Adedirani, M.O. Akando, V.A. Banjoko and G.A. Oluwatosin. 2001. influence of legume fallow on soil properties and yield of maize in south Western Nigeria. *J. Agric. Tropics and Subtropics.* 102(2):109-117.
- Tariq, M. and C.J.B. Mott. 2007. The significance of boron in plant nutrition and environment. *J. Agron.* 6(1):1-10.
- Theocharopoulos, S. P., M. Karayianni, P. Gatzogiani, A. Afentaki and S. Aggelides. 1993. Nitrogen leaching from soils in the Kopais area of Greece. *Soil Use Mgmt.* 9:76-84.
- Threadgill, E. D. 1985. Chemigation via sprinkler irrigation current status and future development. *Appl. Engg. Agric.* 1:16-23.
- Vasane, S.R., P.G. Bhoi, A.S. Patil and A.D. Tumbare. 1996. Effect of liquid fertilizer through drip irrigation on yield and NPK uptake of tomato. *J. Maharashtra Agric. Univ.* 21(3):488-489.
- Vyakaranahal, B.S., M. Shekhargouda, S.A. Patil, A.S. Prabhakar and K. Giriraj. 2001. Effects of planting date and chemicals (TIBA and boron) spray on seed yield, its attributes and quality of single and three-way cross sunflower hybrids during different seasons. *Seed Res.* 29(2):127-135.
- Weil, R.R., R.A. Weismiller and R. S. Turner. 1990. Nitrate contamination of groundwater under irrigated coastal plain soils. *J. Envir. Quality.* 19:441-448.
- Welch, R.M. 1995. Micronutrient nutrition of plants. *Crit. Rev. Plant Sci.* 14:49-82.